




ALLEN & HANBURY'S

Package Leaflet: Information for the User

**Ventolin™ Accuhaler™**  
200 micrograms  
salbutamol sulphate



**Read all of this leaflet carefully before you start taking this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, nurse or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor, nurse or pharmacist.

**In this leaflet:**

- 1 What Ventolin Accuhaler is and what it is used for
- 2 Before you use Ventolin Accuhaler
- 3 How to use Ventolin Accuhaler
- 4 Possible side effects
- 5 How to store Ventolin Accuhaler
- 6 Further information

## 1 What Ventolin Accuhaler is and what it is used for

Ventolin Accuhaler contains a medicine called salbutamol. This belongs to a group of medicines called bronchodilators.

- Bronchodilators help the airways in your lungs to stay open. This makes it easier for air to get in and out.
- They help to relieve chest tightness, wheezing and cough.

Ventolin Accuhaler is used to treat breathing problems in people with asthma and similar conditions. This includes relieving and preventing asthma brought on by exercise or other "triggers". These are things, which bring on asthma symptoms in some people. Common triggers include house dust, pollen, cats, dogs and cigarette smoke.

## 2 Before you use Ventolin Accuhaler

**Do not use Ventolin Accuhaler if:**

- you are allergic (hypersensitive) to salbutamol sulphate or the other ingredient lactose.

**Take special care with Ventolin Accuhaler**

Check with your doctor, nurse or pharmacist before taking your medicine if:

- you have high blood pressure
- you have an overactive thyroid gland
- you have a history of heart problems such as an irregular or fast heartbeat or angina.

## Taking other medicines

Please tell your doctor, nurse or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. This includes herbal medicines. Remember to take this medicine with you if you have to go to hospital.

In particular tell your doctor, nurse or pharmacist if you are taking:

- medicines for an irregular or fast heartbeat
- other medicines for your asthma.

## Using Ventolin Accuhaler with food and drink

You can use Ventolin Accuhaler at any time of day, with or without food.

## Pregnancy and breast-feeding

Talk to your doctor before taking this medicine if you are pregnant, might become pregnant or are breast-feeding.

## Driving and using machines

Ventolin is not likely to affect you being able to drive or use any tools or machines.

## 3 How to use Ventolin Accuhaler

Always use your Ventolin Accuhaler exactly as your doctor has told you. You should check with your doctor, nurse or pharmacist if you are not sure.

## Adults, adolescents aged 12 years and over and children 4 to 11 years of age

- to relieve asthma - One inhalation (200 micrograms).
- to prevent asthma - One inhalation (200 micrograms) 10 to 15 minutes before exercise or exposure to a "trigger".
- the maximum dose is one inhalation (200 micrograms) four times a day.

## Instructions for use


- Your doctor, nurse or pharmacist should show you how to use your inhaler. They should check how you use it from time to time. Not using the Ventolin Accuhaler properly or as prescribed may mean that it will not help your asthma as it should.
- The Ventolin Accuhaler device holds blisters containing Ventolin as a powder
- There is a counter on top of the Ventolin Accuhaler which tells you how many doses are left. It counts down to 0. The numbers 5 to 0 will appear in red to warn you when there are only a few doses left. Once the counter shows 0, your inhaler is empty.

Do not use your inhaler more often than the doctor told you to. Tell your doctor if your medicine does not seem to be working as well as usual, as your chest problem may be getting worse and you may need a different medicine.


Your doctor may have told you to take more than this as an emergency treatment if your wheezing or breathing gets very bad. It is very important that you keep to your doctor's instructions as to how many blisters to take and how often to use your inhaler.

## Using your inhaler

**1** To open your Ventolin Accuhaler, hold the outer case in one hand and put the thumb of your other hand on the thumbgrip. Push your thumb away from you as far as it will go. You will hear a click. This will open a small hole in the mouthpiece.




**2** Hold your Ventolin Accuhaler with the mouthpiece towards you. You can hold it in either your right or left hand. Slide the lever away from you as far as it will go. You will hear a click. This places a dose of your medicine in the mouthpiece.



Every time the lever is pushed back a blister is opened inside and the powder made ready for you to inhale. Do not play with the lever as this opens the blisters and wastes medicine.


**3** Hold the Ventolin Accuhaler away from your mouth, breathe out as far as is comfortable. Do not breathe into your Ventolin Accuhaler. Do not breathe in again yet.

**4** Put the mouthpiece to your lips; breathe in steadily and deeply through the Ventolin Accuhaler with your mouth, not through your nose. Remove the Ventolin Accuhaler from your mouth.



Hold your breath for about 10 seconds or for as long as is comfortable. Breathe out slowly.

**5** To close the Ventolin Accuhaler, slide the thumbgrip back towards you, as far as it will go. You will hear a click. The lever will return to its original position and is reset. Your Accuhaler is now ready for you to use again.



### Cleaning your inhaler

Wipe the mouthpiece of the Accuhaler with a dry tissue to clean it.

### If you use more Ventolin than you should

If you use more than you should, talk to a doctor as soon as possible. The following effects may happen:

- your heart beating faster than usual
- you feel shaky.

These effects usually wear off in a few hours.

### If you forget to use Ventolin Accuhaler

- If you forget a dose, take it as soon as you remember it
- However, if it is time for the next dose, skip the missed dose
- Do not take a double dose to make up for a forgotten dose.

### If you stop using Ventolin Accuhaler

Do not stop using Ventolin Accuhaler without talking to your doctor.

If you have any further questions on the use of this product, ask your doctor, nurse or pharmacist.

## 4 Possible side effects

**If your breathing or wheezing gets worse straight after taking this medicine, stop using it immediately, and tell your doctor as soon as possible.**

Like all medicines, Ventolin Accuhaler can cause side effects, although not everybody gets them. The following side effects may happen with this medicine:

### Allergic Reactions (affects less than 1 in 10,000 people)

If you have an allergic reaction, stop taking Ventolin and see a doctor straight away. Signs of an allergic reaction include: swelling of the face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing, itchy rash, feeling faint and light headed, and collapse.

### Talk to your doctor as soon as possible if:

- you feel your heart is beating faster or stronger than usual (palpitations). This is usually harmless, and usually stops after you have used the medicine for a while
- you may feel your heartbeat is uneven or it gives an extra beat
- these affect less than 1 in 10 people.

If any of these happen to you, talk to your doctor as soon as possible. **Do not stop using this medicine unless told to do so.**

Tell your doctor if you have any of the following side effects which may also happen with this medicine:

### Common (affects less than 1 in 10 people)

- feeling shaky
- headache.

### Uncommon (affects less than 1 in 100 people)

- mouth and throat irritation
- muscle cramps.

### Rare (affects less than 1 in 1,000 people)

- a low level of potassium in your blood
- increased blood flow to your extremities (peripheral dilatation).

### Very rare (affects less than 1 in 10,000 people)

- changes in sleep patterns and changes in behaviour, such as restlessness and excitability.

### The following side effects can also happen but the frequency of these are not known:

- chest pain, due to heart problems such as angina. Tell your doctor, nurse or pharmacist if this occurs. Do not stop using this medicine unless told to do so.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor, nurse or pharmacist.

### If you think this medicine is not working well enough for you

If your medicine does not seem to be working as well as usual, talk to your doctor as soon as possible. Your chest problem may be getting worse and you may need a different medicine. Do not take extra doses of Ventolin Accuhaler unless your doctor tells you to.

## 5 How to store Ventolin Accuhaler

- Keep out of the reach and sight of children.
- Do not store above 30°C.
- Do not use Ventolin Accuhaler after the expiry date, which is stated on the label and carton after 'EXP'. The expiry date refers to the last day of that month.
- If you are told to stop taking this medicine return any unused Ventolin Accuhaler to your pharmacist to be destroyed.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## 6 Further information

### What Ventolin Accuhaler contains

- The active substance is salbutamol sulphate.
- The other ingredient is lactose.
- Each blister contains 200 micrograms of salbutamol sulphate.

### What Ventolin Accuhaler looks like and contents of the pack

Ventolin Accuhaler contains a foil strip. The foil protects the powder for inhalation from the effects of the atmosphere.

Each dose is pre-dispensed.

Each inhaler contains 60 doses of 200 micrograms of salbutamol sulphate.

### Marketing Authorisation holder

Glaxo Wellcome UK Limited trading as:

Allen & Hanburys

Stockley Park West

Uxbridge

Middlesex UB11 1BT

### Manufacturer

Glaxo Wellcome Production

Zone Industrielle No.2

23 Rue Lavoisier

27000 Evreux

France

### Other formats:

To listen to or request a copy of this leaflet in Braille, large print or audio please call, free of charge:

0800 198 5000 (UK only)

Please be ready to give the following information:

**Product name** Ventolin Accuhaler

Reference number 10949/0252

This is a service provided by the Royal National Institute of Blind People.

Leaflet date: August 2011

Ventolin and Accuhaler are trademarks of the GlaxoSmithKline group of companies

© 2011 GlaxoSmithKline group of companies



ALLEN & HANBURYS